



TECH DATA SHEET

REPORTER VIRUS PARTICLES

DESCRIPTION

Product	RVP-1226L, Influenza A Reporter Virus Particles (RVPs)
Lot	IAL-1047A
Subtype	H3N2
Strain	Washington/GKISBBBH95552/2025
Reporter	<i>Renilla</i> Luciferase
Size	1.0 mL/vial
Packaging	10% FBS/DMEM/10mM HEPES
Recommended Input	1 μ L per well (96-well plate) for a S:B >200*
Mycoplasma Test	Negative
Expiration Date	March 2028

SAFETY & HANDLING

Shipping	Shipped on dry ice
Stability and Storage	Store at $\leq -80^{\circ}\text{C}$ upon receipt

* Determined in the MDCK-SIAT1 cell line. Signal-to-background values can vary depending on instrument make, model, and/or calibration.


Influenza A RVPs are used to test the ability of serum, antibodies, and drugs to neutralize infectivity. RVPs display antigenically correct HA/NA protein pseudotyped on replication-incompetent virus particles that contain a heterologous lentiviral (HIV) core. RVPs are capable of a single round of infection and carry a genome that expresses either a GFP or luciferase optical reporter gene upon infection. RVPs are produced in HEK-293T cells using four separate plasmids, encoding the HA protein, the NA protein, a lentiviral gag polyprotein, and a reporter gene.

RVPs are created using a second-generation lentiviral system with components that are highly unlikely to recombine to produce a fully infectious virus (requiring 3 separate recombination events to do so). However, lentiviruses are capable of genomic integration and RVPs are derived from biological materials so should be handled with caution within a BSL2 or enhanced BSL2 laboratory environment. RVPs are not to be used in humans or in animals raised for food.

Thaw tubes in a 37°C water bath for 3 minutes and place on ice until ready to use. RVPs will appear as a translucent solution. Gently mix prior to use and pulse tube for 3 seconds at high speed in a tabletop microfuge to recover all volume from the tube. Vortexing of RVPs should be avoided. Re-freezing of RVPs is not recommended.

INFECTIVITY DATA

Influenza A RVPs (Lot IAL-1047A)



Infectivity is determined in MDCK-SIAT1 cells. Infectivity data represents the average of three independent vials, each tested in quadruplicate.

Renilla luciferase activity measured using the Promega *Renilla*-Glo luciferase assay system (Promega #E2710). Sample luminescence was read using a Perkin-Elmer Envision plate reader.

SIGNAL TO BACKGROUND

RVP 384-well Input (μl)	Signal: Background
20	15642
10	4981
5	2318
2.5	1775
1.25	1747
0.63	683
0.31	396

Signal to background is calculated using mock infected cells as the negative control value.